







The reappearance of *Puma concolor* (Linnaeus, 1771) (Mammalia, Carnivora, Felidae) in the city of Rio de Janeiro, Brazil

Jorge Antônio Lourenço Pontes^{1*}, Rafael Andrada de Araújo Martins², Luiz Eduardo Mendonça Regio¹, Mário Luiz Gomes Soares³, Filipe de Oliveira Chaves³, Helena Godoy Bergallo⁴

1 Programa de Pós-Graduação em Ensino de Ciências, Ambiente e Sociedade Universidade do Estado do Rio de Janeiro, São Gonçalo, RJ, Brazil • JP pontesjal@hotmail.com  <https://orcid.org/0000-0003-2741-4273> • LR luizregio@hotmail.com  <https://orcid.org/0000-0002-6588-1123>

2 Laboratório de Zoologia de Vertebrados, Centro Universitário São José, Rio de Janeiro, RJ, Brazil • RA andrada88rafael@gmail.com  <https://orcid.org/0000-0001-7663-8949>

3 Núcleo de Estudos em Manguezais, Faculdade de Oceanografia, Universidade do Estado do Rio de Janeiro, Rio de Janeiro, RJ, Brazil • MS mariolgs@gmail.com  <https://orcid.org/0000-0002-3312-7257> • FC fochaves@gmail.com  <https://orcid.org/0000-0001-7114-7585>

4 Departamento de Ecologia, Universidade do Estado do Rio de Janeiro, Rio de Janeiro, RJ, Brazil • HB nena.bergallo@gmail.com  <https://orcid.org/0000-0001-9771-965X>

* Corresponding author

Abstract

Large mammals, especially felids such as Mountain Lion, *Puma concolor* (Linnaeus, 1771), are disappearing from their original habitats, due to the loss of natural areas and hunting, especially in the metropolitan regions. In the city of Rio de Janeiro, Brazil, *P. concolor* had not been observed for almost a century, despite the city having forest fragments as large as 10,000 ha. Here we present records that confirm the reappearance of this species in the city of Rio de Janeiro, where it has been considered extirpated.

Keywords

Atlantic Forest, extirpation, metropolitan region, protected areas, wild cats

Academic editor: Guilherme S. T. Garbino | Received 14 June 2021 | Accepted 31 August 2021 | Published 28 September 2021

Citation: Pontes JAL, Martins RAA, Regio LEM, Soares MLG, Chaves FO, Bergallo HG (2021) The reappearance of *Puma concolor* (Linnaeus, 1771) (Mammalia, Carnivora, Felidae) in the city of Rio de Janeiro, Brazil. Check List 17 (5): 1353–1358. <https://doi.org/10.15560/17.5.1353>

Introduction

Mammal populations, especially of large predators, are declining in the city of Rio de Janeiro, with some species such as the Jaguar, *Panthera onca* (Linnaeus, 1758), already extirpated (Vaz 1985; Bergallo et al. 2000; Vera Y Conde et al. 2000). The Mountain Lion, *Puma concolor* (Linnaeus, 1771), has a geographic distribution that encompasses both North and South America, from Canada to the Patagonian Argentina (Oliveira and Cassaro 2005; Reis et al. 2011; Nielsen et al. 2015). Six

subspecies of *P. concolor* are recognized (Culver et al. 2000), this species not considered globally threatened and is classified as Least Concern in the IUCN Red List of Threatened Species (Nielsen et al. 2015), but the population from eastern United States, was extirpated in 2018 (Racey and Kloer 2018). Despite the broad distribution in all Brazilian biomes, populations of *P. concolor* subspecies are in decline in several regions, such as in the Caatinga (*P. concolor greeni* (Nelson & Goldman, 1931))

and in the south-southeastern Atlantic Forest (*P. concolor concolor* (Linnaeus, 1758)) (Reis et al. 2011; Matte 2012; Nascimento and Garbino 2013; Cronemberger et al. 2018). *Puma concolor* is classified as Vulnerable in Brazil (Brasil 2014) and the state Rio de Janeiro (Estado do Rio de Janeiro 1998; Bergallo et al. 2000), mainly due to habitat loss and poaching (Rocha et al. 2003; Bergallo et al. 2009).

There were no records proving the occurrence of *P. concolor* for more than 80 years from the city of Rio de Janeiro (Côrrea 2017), despite having been considered a rare species in the 1980s, without evidence (Vaz 1985). To the best of our knowledge, it has not been documented in Rio de Janeiro in the last 80 years (Corrêa 2017), and it was officially declared Extinct in the Municipal List of Threatened Species of Flora and Fauna (Vera Y Conde et al. 2000).

Despite anecdotal comments on the reappearance of *P. concolor* in Rio de Janeiro, in the two largest forest fragments on the Gericinó-Mendanha and Pedra Branca mountain ranges, no unambiguous records (e.g., photographs, footprints) have been available until now. Here, we confirm the current presence of *P. concolor* in the city of Rio de Janeiro.

Study Area

Pedra Branca State Park (PE Pedra Branca) covers an

area of 12,492 ha and is inserted in the Pedra Branca Mountain Range in the city of Rio de Janeiro (Rio de Janeiro hereafter) (Coura et al. 2009; INEA 2013; Castro 2015) (Fig. 1).

The Reserva Particular do Patrimônio Natural Bicho Preguiça (RPPN Bicho Preguiça hereafter) has an area of 1.73 ha and is located in the Gericinó-Mendanha mountain range. Mendanha State Park (PE Mendanha) and Serra do Mendanha Natural Municipal Park (PNM Serra do Mendanha) are also located in the same mountain range (INEA 2014; Castro 2015; Martins and Pontes 2019) (Fig. 1).

The Guaratiba State Biological Reserve (RBEG) is approximately 50 km from downtown Rio de Janeiro, located on the Sepetiba/Guaratiba coastal plain in the Piracão and Portinho watersheds. It covers approximately 3,360 ha along the eastern side of Sepetiba Bay, between the Piraquê River and Guaratiba Hill (Fig. 1). The reserve encompasses a complex of islands, rivers, and canals with extensive mangrove forests (2,090 ha) and salt flats (740 ha) (Estrada et al. 2013; Castro 2015; Soares et al. 2017).

RBEG is inserted in a highly connected landscape characterized by several subdivisions: (i) Pedra Branca mountain range; (ii) Coastal plain (rivers, freshwater wetlands, coastal scrub (restinga hereafter), salt-flats and mangrove forests); (iii) Sepetiba Bay (estuary, tidal




Figure 1. Map *Puma concolor* records in the city of Rio de Janeiro: 1 and 2 = RBEG Guaratiba (2007 and 2008); 3 = PE Mendanha and PNM Serra do Mendanha (2008); 4 = RPPN Bicho Preguiça (2018 and 2020); 5 = Roberto Burle Marx Property, Barra de Guaratiba, contiguous to PE Pedra Branca (2020). Insert maps show the location of Rio de Janeiro state in southeastern Brazil.

channels, mangrove forests on islands and mud banks); (iv) Restinga da Marambaia (mangrove forests, restinga, dunes, freshwater wetlands, and beaches); and (v) Ocean.

The connection between the types of ecosystems in this region can also be observed, not only by differences in vegetation, but also by the flow of freshwater from the slopes of the Pedra Branca mountain range to the coastal plain, which feeds the shallow (1.5 m) Guaratiba aquifer (Vicent et al. 2010).

Methods

An inventory of terrestrial mammals has been ongoing since June 2015 in the RPPN Bicho Preguiça. Six camera traps were set up without the use of baits, combined with 1,540 h of active search on trails where *Puma concolor* footprints were observed (Martins and Pontes 2019). Field equipment included a laptop to download data, cell phones with a georeferencing app, camera traps, gypsum to make footprint molds, and a ruler in millimeters.

Geographic coordinates use the SIRGAS 2000 datum.

Additionally, records derived from other Universidade do Estado do Rio de Janeiro field activities were incorporated in our study: the herpetofauna of Serra do Mendanha by the Laboratório de Ecologia de Vertebrados, and mangrove forests and salt flats in the RBEG by the Núcleo de Estudos em Manguezais.

Results

Puma concolor (Linnaeus, 1771)

Figures 1, 2

New records. BRAZIL – Rio de Janeiro • city of Rio de Janeiro; near the Piracão River RBE Guaratiba; 23.01°S, 043.57°W; 2 m alt.; Bergallo and Pontes leg.; 2007; footprint • city of Rio de Janeiro; RBE Guaratiba; salt flats associated with mangrove forests; 23.01°S, 043.60°W; 2 m alt.; Soares and collaborators leg.; 2008; footprint and




Figure 2. Evidence of *Puma concolor* in the city of Rio de Janeiro. **A.** Scratch marks on a tree trunk in RPPN Bicho Preguiça, Gericinó-Mendanha massif (2020). **B.** Footprint mold of anterior paw in RPPN Bicho Preguiça, Gericinó-Mendanha mountain range (2019). **C.** Anterior paw footprint recorded at 700 m altitude in PNM Serra do Mendanha, Gericinó-Mendanha mountain range (2008). **D.** Individual at the gate of the Roberto Burle Marx Property, Barra de Guaratiba, Pedra Branca mountain range (June 2020). **E, F.** Footprints RBE Guaratiba, Barra de Guaratiba (2007 and 2008). Photographs by Rafael Andrada (A, B); Jorge Pontes (C); security camera in the Roberto Burle Marx Property (D); Núcleo de Estudos em Manguezais (E, F).

scratch marks on tree trunks • city of Rio de Janeiro; PNM Serra do Mendanha; 22.82°S, 043.50°W; 700 m alt.; Pontes and collaborators; 05.VII.2008; footprint • city of Rio de Janeiro; RPPN Bicho Preguiça; 22.77°S, 043.60°W; 245 m alt.; Martins and Pontes leg.; 16.VIII.2019; footprint and mold • city of Rio de Janeiro; RPPN Bicho Preguiça; 22.83°S, 043.55°W; 321 m alt.; Martins and Pontes leg.; 15.VIII.2020; scratch marks on tree trunks • city of Rio de Janeiro; Roberto Burle Marx Property, Pedra Branca mountain range; 23.02°S, 043.55°W; 15 m alt.; Pontes leg.; 30.VI.2020; security camera.

Identification. *Puma concolor* is a large cat of the sub-family Felinae and the second-largest New World cat after Jaguar, *Panthera onca* (Linnaeus, 1748), from which it can be differentiated by the lack of spots and a uniform color dorsal pelage. *Puma concolor* can reach 230 cm in length from snout to tail and more than 70 kg in body mass. Its pelage varies in color from lighter grayish brown to darker reddish brown, and the tip of the tail is black. Another species of cat, *Herpailurus yagouaroundi* (É. Geoffroy Saint-Hilaire, 1803), can occur in anthropic environments (Culver et al. 2000; Leite 2000; Matte 2012; Caso et al. 2015; Ferreira et al. 2018; ICMBio 2018; Martins and Pontes 2019), but *H. yagouaroundi* can be differentiated from *P. concolor* by its smaller size (approximately 8 kg and 80 cm total length), more elongated body, short legs, and dark brown color, as well as not being from a genetically close group (Culver et al. 2000; Caso et al. 2015; Reis et al. 2011). Secretive and largely solitary, *P. concolor* is both nocturnal and crepuscular, although daytime sightings do occur. This species is an ambush predator which pursues a wide variety of prey. Individuals are territorial and live at low population densities (Silva 1994; Oliveira and Cassaro 2005; Reis et al. 2011; Matte 2012; ICMBio 2018).

Discussion

Our new data confirm that *Puma concolor* is not extirpated in coastal Rio de Janeiro state and that it is recolonizing the city of Rio de Janeiro. It is now present at least in the Gericinó-Mendanha and Pedra Branca mountain ranges and in the RBEG (Figs. 1, 2).

Individuals of *P. concolor* in the city of Rio de Janeiro may have dispersed from the Serra do Tinguá and surrounding areas (Cronemberger et al. 2018), first colonizing the Gericinó-Mendanha mountain range, then the Pedra Branca mountain range and lowlands, as in the RBEG. The presence of Collared Pecary, *Dicotyles tajacu* (Linnaeus, 1758), in the Gericinó-Mendanha mountain range (Martins and Pontes 2020), may attract and provide prey for *P. concolor*. Wild pigs are typically prey of *P. concolor* (Sana and Cullen Jr 2008; Reis et al. 2011).

Although *P. concolor* is not usually hunted in Rio de Janeiro state (Ferreira et al. 2018), there was a recent social media post of a mountain lion killed by a hunter

in Serra dos Órgãos (O Globo 2019). Events like this threatened the existence of animals occurring in the fragmented Atlantic Forest (Cullen Jr. et al. 2001; Souza and Alves 2014; Galetti et al. 2016; Cronemberger et al. 2018), especially in densely populated areas such as the Rio de Janeiro metropolitan region, where hunting is still a common, albeit illegal, practice (Vaz 1985; Ferreira et al. 2018).

The record of *P. concolor* made by a security camera in the Roberto Burle Marx Property, which is in the foothills of the Pedra Branca mountain range, and records of footprints on the salt flats in RBEG strongly suggest expanded home range that includes resources available on the Guaratiba coastal plain.

Although the occurrence of wild cats in mangroves is known worldwide (Saenger et al. 1983; Fernandes 2000), there are few records of *P. concolor* in mangroves in Brazil. Mazzolli (1993) reported the occurrence of this species in restinga and mangroves on the coastal plain of Santa Catarina, and according to this author, only one of 22 mountain lion records in the state was obtained on the coastal plain of that state, probably due to the higher human population along the coast. As in northern Santa Catarina, coastal areas of Paraná and southern São Paulo states are more sparsely populated (Mazzolli 1993). In fact, Leite (2000) reported the presence of *P. concolor* in Superagui National Park and in the Guaraqueçaba Environmental Protection Area, both in coastal Paraná state. Martins et al. (2008) also reported the occurrence of *P. concolor* on the coast of São Paulo state, in the Juréia-Itatins Ecological Station, although they did not specifically mention that animals were living in mangroves.

In addition to written reports on the presence of wild cats in mangroves, oral communications about observations of animals and footprints in mangroves have been made by members of traditional communities, fishermen, and managers of protected areas; some sightings of animals have been video edition or photographed (Fundação Florestal 2019; G1 2020). Researchers at the Núcleo de Estudos em Manguezais (Universidade do Estado do Rio de Janeiro) have also reported the presence of wild cats in mangroves in Cananéia, Bertioga (both in São Paulo state), Guaratiba (Rio de Janeiro state), and Sucuriju (Amapá state). All these records are from areas where mangroves occur at the border of relatively well-conserved terrestrial ecosystems in landscapes where terrestrial and coastal-estuarine systems are widely connected.

Hutchings and Saenger (1987) highlighted that most terrestrial vertebrates use mangroves as an extension of their native ranges, as alternative refuges, connection between two terrestrial ecosystems, protection against predators, and feeding grounds. Considering that the Guaratiba mangroves are on the border of the coastal landscape and therefore not connecting different terrestrial ecosystems, wild cats in Brazil are occasional mangrove users, and wild cats are at the top of the food chain, one of our hypotheses is use the Guaratiba mangroves to forage for animal prey and refuge.

The occurrence of *P. concolor* in the city of Rio de Janeiro municipality highlights both the importance of conserving forest remnants and the need to maintain and increase connectivity between these remnants. Our records from the Guaratiba lowlands (Baixada de Guaratiba) also suggest the importance of landscape connections between different ecosystem types in the Atlantic Forest domain (Marone et al. 2009).

The conservation of *P. concolor* and other threatened mammal species in the city of Rio de Janeiro depends on state and municipal public policies targeting these species, including proper management and vigilance, especially in the PE Mendanha (4,398 ha) and PE Pedra Branca (12,492 ha) (Castro 2015). However, the protection of smaller reserve areas, both municipally and privately owned (such as RPPN Bicho Preguiça), as well as unprotected natural areas, such as Camboatá Forest, is still important. Such smaller areas can function as steppingstones for the transit of *P. concolor* and other species (Oliveira et al. 2015; Pontes et al. 2015; Pontes et al. 2020).

Acknowledgements

We thank Tadeu de Oliveira for confirming the identification of footprints. HGB thanks Prociência – UERJ, FAPERJ (E-26/202.757/2017) and CNPq (306585/2018-9) for scholarships and support. We thank the Municipal Secretary of Environment of Rio de Janeiro and the State Institute of Environment of Rio de Janeiro for research permits issued to JALP (nº. 002-2010 INEA and 011-2020 SMAC). RAAM thanks Sérgio and Patrícia Martins for funding the Inventory and Monitoring of Terrestrial Mammals in the RPPN Bicho Preguiça, and his son Bernardo for his love and sweetness.

Authors' Contributions

Conceptualization: JALP. Data curation: FOC. Formal analysis: JALP, HGB. Investigation: JALP, RAAM, LEMR, MLGS, FOC, HGB. Methodology: RAAM, MLGS, FOC. Resources: RAAM. Supervision: JALP. Visualization: RAAM, LEMR, FOC. Writing – original draft: JALP, MLGS, HGB. Writing – review and editing: JALP, HGB.

References

- Bergallo HG, Esbérard CEL, Geise L, Grelle CEV, Vieira MV, Gonçalves PR, Paglia A, Attias N (2009) Mamíferos endêmicos e ameaçados no estado do Rio de Janeiro: diagnóstico e estratégias para a conservação. In: Bergallo HG, Fidalgo ECC, Rocha CFD, Uzêda MC, Costa MB, Alves MA, Van Sluys M, Santos MA, Costa TCC, Cozzolino AC (Org.) Estratégias e ações para a conservação da biodiversidade no estado do Rio de Janeiro. Instituto Biomas, Rio de Janeiro, Brazil, 209–219.
- Bergallo HG, Geise L, Bonvicino CR, Cerqueira R, D'Andrea OS, Esbérard CEL, Fernandez FAZ, Grelle CE, Peracchi A, Siciliano S, Vaz SM (2000) Mamíferos. In: Bergallo HG, Rocha CFD, Alves MAS, Van Sluys M (Eds.) A fauna ameaçada de extinção do Estado do Rio de Janeiro. EdUERJ, Rio de Janeiro, Brazil, 125–135.
- Brasil (2014) Portaria MMA nº 444 de 17 de dezembro de 2014. Diário Oficial da União, Brasília, Brazil 245: 121–126.
- Caso A, de Oliveira T, Carvajal SV (2015) *Herpailurus yagouaroundi*. The IUCN Red List of Threatened Species 2015: e.T9948A50653167. <https://doi.org/10.2305/iucn.uk.2015-2.rlts.t9948a50653167.en>
- Castro PF (Ed.) (2015) Atlas das unidades de conservação do estado do Rio de Janeiro (2ª edição.). Metalivros, São Paulo, Brazil, 172 pp.
- Cheida CC, Rodrigues FHG (2014) Introdução a técnicas de estudo em campo para mamíferos carnívoros terrestres. In: Reis NR, Peracchi AL, Rossaneis BK, Fregonezi MN (Eds.) Técnicas de estudos aplicadas aos mamíferos silvestres brasileiros, 2ª edição. Technical Books Editora, Rio de Janeiro, Brazil, 105–136.
- Corrêa MA (2017) O sertão carioca, 2ª edição. Contra Capa, Rio de Janeiro, Brazil, 328 pp.
- Coura PHF, Souza GM, Fernandes MC (2009) Mapeamento geológico da susceptibilidade à ocorrência de incêndios no maciço da Pedra Branca, município do Rio de Janeiro. Anuário do Instituto de Geociências 32 (2): 14–25.
- Cronemberger C, Pereira, FA, Bacellar AEF, Silva LG (2018) First record of leucism in puma from Serra dos Órgãos National Park, Brazil. CATnews 68: 38–41. <https://orcid.org/0000-0002-7993-9015>
- Cullen Jr L, Bodmer ER, Valladares-Padua C (2001) Ecological consequences of hunting in Atlantic Forest patches, São Paulo, Brazil. Oryx 35 (2): 137–144. <https://doi.org/10.1046/j.1365-3008.2001.00163.x>
- Culver M, Johnson WE, Pecon-Slattery J, O'Brien SJ (2000) Genomic ancestry of the American puma (*Puma concolor*). The Journal of Heredity 91 (3): 186–197. <https://doi.org/10.1093/jhered/91.3.186>
- Estado do Rio de Janeiro (1998) Portaria SEMA 01 de 05 de junho de 1998. Lista das espécies da fauna ameaçadas de extinção no estado do Rio de Janeiro. Diário Oficial do Estado do Rio de Janeiro 102: 9–16.
- Estrada GCD, Soares MLG, Chaves FO, Cavalcanti VF (2013) Analysis of the structural variability of mangrove forests through the physiographic types approach. Aquatic Botany 111: 135–142. <https://doi.org/10.1016/j.aquabot.2013.06.003>
- Fernandes MEB (2000) Association of mammals with mangrove forests: a worldwide review. Boletim do Laboratório de Hidrobiologia 13: 83–108.
- Ferreira ET, Albuquerque SVP, Aximoff IA, Marins JRGA, Pontes JAL (2018) Aves e mamíferos cinegéticos nas unidades de conservação da natureza de proteção integral do estado do Rio de Janeiro. Revista Ineana 6 (3): 38–53.
- Fundação Florestal (2019) Onça-parda é avistada no Parque Estadual Restinga de Bertiooga. <https://www.infraestruturameioambiente.sp.gov.br/fundacaoflorestal/2019/11/onca-parda-e-avistada-no-parque-estadual-restinga-de-bertiooga/>. Accessed on: 2021-2-08.
- G1 (2020) Onça-parda é flagrada atravessando canal no litoral de SP. <https://g1.globo.com/sp/santos-regiao/noticia/2020/07/28/onca-parda-e-flagrada-atravessando-canal-no-litoral-de-sp-video.ghtml>. Accessed on: 2021-2-08.
- Galetti M Brocardo, CR, Begotti RA, Hortenci L, Rocha-Mendes F, Bernardo CSS, Bueno RS, Nobre R, Bovendorp RS, Marques RM, Meirelles F, Gobbo SK, Beca G, Schmaedecke G, Siqueira T (2016) Defaunation and biomass collapse of mammals in the largest Atlantic Forest remnant. Animal Conservation 20 (3): 270–281 <https://doi.org/10.1111/acv.12311>
- Hutchings PA, Saenger P (1987) Ecology of mangroves. University of Queensland Press, Brisbane, Australia, 388 pp.
- ICMBio (Instituto Chico Mendes de Conservação da Biodiversidade) (2018) Sumário executivo do plano de ação nacional para a conservação da onça-parda. <https://www.icmbio.gov.br/portal/faunabrasileira/plano-de-acao-nacional-lista/2466-plano-de-acao-nacional-para-conservacao-da-onca-parda>. Accessed on: 2021-7-8.
- INEA (Instituto Estadual do Ambiente) (2013) Plano de manejo do

- Parque Estadual da Pedra Branca, Resolução INEA Nº 74 de 02 de Julho de 2013. <http://www.femerj.org/wp-content/uploads/Plano-de-manejo-do-Parque-Estadual-da-Pedra-Branca-PEPB-2.pdf>. Accessed on: 2021-2-08.
- Leite MRP (2000) Relações entre a onça-pintada, onça-parda e moradores locais em três unidades de conservação da Floresta Atlântica do Estado do Paraná, Brasil. Master's thesis, Universidade Federal do Paraná, Curitiba, Brazil, 75 pp.
- Marone E, Soares MLG, Creed JC, Zalmom IR, Villaça RC, Chaves LCT, Rêgo PJA (2009) Ecossistemas marinhos. In: Bergallo HG, Fidalgo ECC, Rocha CFD, Uzêda MC, Costa MB, Alves MA, Van Sluys M, Santos MA, Costa TCC, Cozzolino AC (Eds.) Estratégias e ações para a conservação da biodiversidade no estado do Rio de Janeiro. Instituto Biomas, Rio de Janeiro, Brazil, 235–245.
- Martins RAA, Pontes JAL (2019) Inventário e monitoramento da mastofauna terrestre da Reserva Particular do Patrimônio Natural Bicho Preguiça, Rio de Janeiro, Brasil: dados preliminares. In: Anais 1º, BIOEMFOCO. Centro Universitário São José Rio de Janeiro, Brazil, 1-2. <https://www.researchgate.net/publication/341998625>. Accessed on: 2020-11-03.
- Martins RAA, Pontes JAL (2020) Registro da ocorrência de *Pecari tajacu* (Linnaeus, 1758), uma espécie que era declarada extinta no município do Rio de Janeiro, estado do Rio de Janeiro, Sudeste do Brasil. Boletim Sociedade Brasileira de Mastozoologia 88: 58–61. <https://doi.org/10.13140/rg.2.2.21198.08007>
- Martins R, Quadros J, Mazzolli M (2008) Hábito alimentar e interferência antrópica na atividade de marcação territorial do *Puma concolor* e *Leopardus pardalis* (Carnivora: Felidae) e outros carnívoros na Estação Ecológica de Juréia-Itatins, São Paulo, Brasil. Revista Brasileira de Zoologia 25 (3): 427–435. <https://doi.org/10.1590/s0101-81752008000300007>
- Matte EM (2012) Filogeografia de *Puma concolor* (Carnivora, Felidae) na América do Sul. PhD thesis, Universidade Federal do Rio Grande do Sul, Porto Alegre, Brazil, 109 pp.
- Mazzolli M (1993) Ocorrência de *Puma concolor* (Linnaeus) (Felidae, Carnivora) em áreas de vegetação remanescente de Santa Catarina, Brasil. Revista Brasileira de Zoologia 10 (4): 581–587. <https://doi.org/10.1590/s0101-81751993000400002>
- Nascimento FO, Garbino GST (2013) On the available names for the northern and eastern South American subspecies of *Puma concolor* (Linnaeus, 1771) (Mammalia: Felidae) and their type localities. Zootaxa 3646 (2): 194–198. <https://doi.org/10.11646/zootaxa.3646.2.8>
- Nielsen C, Thompson D, Kelly M, Lopez-Gonzalez CA (2015) *Puma concolor* (errata version published in 2016). The IUCN Red List of Threatened Species 2015: e.T18868A97216466. <https://doi.org/10.2305/iucn.uk.2015-4.rlts.t18868a50663436.en>
- O Globo (2019) Coluna Ancelmo Gois. O Globo (newspaper) Rio de Janeiro, Brazil, 2019-19-04, Matutina, Rio: 12. <https://acervo.oglobo.globo.com/consulta-ao-acervo/?navegacaoPorData=201020190418>. Accessed on: 2020-11-03.
- Oliveira DR, Paes-Pinto MP, Silva LMR, Nunes RP, Pontes JAL (2015) Os Mamíferos terrestres do Parque Natural Municipal Bosque da Barra, Barra da Tijuca, Rio de Janeiro. In: Pontes JAL (Ed.) Biodiversidade carioca: segredos revelados. Technical Books, Rio de Janeiro, Brazil, 246–256.
- Pontes JAL, Bergallo HG, Lira FO, Alves MAS (2020) Parecer e avaliação do estudo de impacto ambiental e da audiência pública sobre a construção do autódromo internacional na Floresta do Camboatá com ênfase na fauna. Deodoro, Rio de Janeiro, estado do Rio de Janeiro. SOS Floresta do Camboatá, Rio de Janeiro, Brazil. <https://doi.org/10.13140/rg.2.2.17279.02729>
- Pontes JAL, Pontes RC, Rocha CFD (2009) The snake community of Serra do Mendanha, in Rio de Janeiro state, southeastern Brazil: composition, abundance, richness and diversity in areas with different conservation degrees. Brazilian Journal of Biology 3 (69): 795–804. <https://doi.org/10.1590/s1519-69842009000400006>
- Pontes JAL, Pontes RC, Rocha RF, Lindenberg PM, Silva KP, Santos WA, Lemos NA, Hassan PGA, Alves AO, Lopes LFBA, Perro LCT, Boldrini AP, Nunes ECF, Costa LF, Kisling RW, Rocha CFD (2015) Unidades de conservação da cidade do Rio de Janeiro: hotspots da herpetofauna carioca. In: Pontes JAL (Ed.) Biodiversidade carioca: segredos revelados. Technical Books, Rio de Janeiro, Brazil, 176–194.
- Racey M, Kloer P (2018) Long extinct eastern cougar to be removed from endangered species list correcting Linger anomaly. News Bulletin, U.S. Fish and Wildlife Service. https://www.fws.gov/northeast/ecougar/pdf/Cougar_News_Bulletin_Final_1_18.pdf. Accessed on: 2020-7-02.
- Reis NR, Peracchi AL, Pedro WA, Lima IP (Eds.) (2011) Mamíferos do Brasil, 2ª. edição. Nelio R. dos Reis, Londrina, Brazil, 439 pp.
- Reis NR, Ramos PHG, Andrade FR, Peracchi AL (2014) Técnicas de estudo de mamíferos de médio e grande porte e para os não voadores, de pequeno porte. In: Reis NR, Peracchi AL, Rossaneis BK, Fregonezi MN (Eds.) Técnicas de estudos aplicadas aos mamíferos silvestres brasileiros, 2ª edição. Technical Books Editora, Rio de Janeiro, Brazil, 257–269.
- Rocha CFD, Bergallo HG, Alves MAS, Van Sluys M (2003) A biodiversidade nos grandes remanescentes florestais do Estado do Rio de Janeiro e nas restingas da Mata Atlântica. RiMa Editora, São Carlos, Brazil, 146 pp.
- Saenger P, Hegerl EJ, Davie JDS (Ed.) (1983) Global status of mangrove ecosystems. Working Group on Mangrove Ecosystems of the IUCN, Gland, Switzerland. Commission on Ecology Papers 3: 1–92.
- Sana DA, Cullen Jr. L (2008) *Puma concolor capricornensis* – mamíferos. In: Machado ABM, Drumond GM, Paglia AP (Eds.) Livro Vermelho da fauna brasileira ameaçada de extinção. Ministério do Meio Ambiente, Brasília/Fundação Biodiversitas, Belo Horizonte, Brazil, 795–799.
- Silva F (1994) Mamíferos silvestres – Rio Grande do Sul, 2ª edição. Fundação Zoobotânica do Rio Grande do Sul, Porto Alegre, Brazil, 246 pp.
- Soares MLG, Chaves FO, Estrada GCD, Farnedez V (2017) Mangrove forests associated with salt flats: a case study from southeast Brazil. Brazilian Journal of Oceanography 65 (2): 102–115. <https://doi.org/10.1590/s1679-87592017083006502>
- Souza JB, Alves RRN (2014) Hunting and wildlife use in an Atlantic Forest remnant of northeastern Brazil. Tropical Conservation Science 7 (1): 145–160. <https://doi.org/10.1177/194008291400700105>
- Vaz SM (1985) Mamíferos do Rio de Janeiro. Boletim FBCN 20: 80–89.
- Vera Y Conde CF, Bueno C, Bonvicino CR, Vaz SM, Esberárd CEL (2000) Mamíferos. In: Di Maio FR, Silva MBR (Eds.) Espécies ameaçadas de extinção no Município do Rio de Janeiro, flora e fauna. Secretaria Municipal de Meio Ambiente do Rio de Janeiro, Rio de Janeiro, Brazil, 39–43.
- Wilson DE, Reeder DM (Ed.) (2005) Mammal species of the world. A taxonomic and geographic reference. 3rd edition. Johns Hopkins University Press, Baltimore, USA, 2142 pp.